

Хронический некалькулезный холецистит с билиарным сладжем

Хронический некалькулезный холецистит с билиарным сладжем – это воспалительное заболевание, вызывающее поражение стенки желчного пузыря и моторно-тонические нарушения билиарной системы, сопровождающееся появлением гиперэхогенных частиц в полости желчного пузыря и проявляется болью билиарного типа (1-10). Причиной моторной дисфункции желчного пузыря может выступать сужение пузырного протока, мышечная гипертрофия, хроническое воспаление стенки желчного пузыря.

Термином “билиарный сладж” обозначают любую неоднородность желчи, выявляемую при эхографическом исследовании (2-10).

Билиарный сладж состоит из гранул билирубината кальция, кристаллов моногидрата холестерина и полимеризованных нитей гликопротеинового муцина желчного пузыря (2-10). Ранее в экспериментах *in vitro* было показано, что только кристаллы моногидрата холестерина обладают повышенной эхогенностью без акустической тени при ультразвуковом сканировании (2-5).

Билиарный сладж часто образуется у беременных в третьем триместре, у больных с ожирением, находящихся на низкокалорийной диете, у пациентов, находящихся на парентеральном питании и после гастрэктомии или колонэктомии (11-12).

По расположению в желчном пузыре и форме **билиарный сладж** классифицируют как рассеянный, туморообразный, поверхностный и преципитирующий (5). Показана динамика его трансформации в холестериновые желчные камни: рассеянный билиарный сладж → поверхностный билиарный сладж → преципитирующий билиарный сладж → холестериновый желчный камень без акустической тени → холестериновый желчный камень с акустической тенью (5). Время необходимое для этого составляет от 3 до 36 месяцев (12). Процент трансформации колеблется от 5% до 50% в зависимости от причины (12). Так, у 25-50% больных ожирением, находящихся на низкокалорийной диете в течение 3-6 месяцев, формируется билиарный сладж и холестериновые желчные камни (12). У 40% больных, страдающих ожирением – через 6 месяцев после операции на желудке (12).

Ежедневный прием 600 мг УДХК снижает риск образования холестериновых желчных камней с 28% до 3% у этих пациентов (12). У 45% взрослых и у 43% больных детей, находящиеся в течение 3-4 месяцев на парентеральном питании, образуются холестериновые желчные камни в желчном пузыре (12). Во время третьего триместра беременности у 30% беременных женщин формируется билиарный сладж, у 2% – холестериновые желчные камни в желчном пузыре (12). После родов моторная функция желчного пузыря восстанавливается и билиарный сладж исчезает у 60-70% рожениц, а холестериновые желчные камни спонтанно растворяются у 20-30% (12).

Российские гастроэнтерологи выделяют 3 основных типа билиарного сладжа, имеющих наиболее четко очерченную эхографическую картину (11):

1. **Микролитиаз** – взвесь гиперэхогенных частиц в виде точечных, единичных или множественных, смещаемых гиперэхогенных образований, не дающих акустической тени, выявляемых после изменения положения тела пациента.
2. **Эхонеоднородная желчь** с наличием различной плотности сгустков, смещаемых и не дающих акустической тени, или в редких случаях с эффектом ослабления за сгустком.
3. **Сочетание замазкообразной желчи с микролитами.** При этом микролиты могут быть одновременно как в составе сгустка замазкообразной желчи, так и в полости желчного пузыря.

Диагностические критерии хронического некалькулезного холецистита с билиарным сладжем

1. Боли и ощущение дискомфорта в области правого подреберья, постоянные или возникающие периодически, различной интенсивности и продолжительности несвязанные или связанные с приемом пищи (1).

Кроме того, боль может сочетаться с одним и более из следующих признаков:

- a. Постоянное или периодически возникающее ощущение горечи во рту
- b. Тошнота, изжога, отрыжка
- c. Постоянное или периодически возникающее ощущение вздутия и урчания в животе
- d. Неустойчивый стул с преобладанием запоров или поносов
- e. Возникновение боли в ночное время
2. **Нарушение эвакуаторной функции желчного пузыря.**
3. По данным УЗИ исследования **утолщение стенки желчного пузыря до 3-6 мм.**

Причины болевого синдрома, хронического воспаления в стенке и нарушения эвакуаторной функции желчного пузыря

1. Патология гладкомышечных и эпителиальных клеток желчного пузыря (**повышенная экспрессия ЦОГ-2 в гладкомышечных и эпителиальных клетках стенки желчного пузыря**).
2. Гиперсекреция гликопротеинового муцина в просвет желчного пузыря и повышение концентрации гликопротеинового муцина в пузирной желчи свыше точки полимеризации (>2.0 мг/мл) (**избыточная экспрессия ЦОГ-2 в эпителиальных клетках стенки желчного пузыря**).
3. Дискоординация желчного пузыря и пузирного протока (**повышенная экспрессия ЦОГ-2 в гладкомышечных клетках желчного пузыря и пузирного протока**).
4. Увеличение сопротивления пузирного протока (**повышенная экспрессия ЦОГ-2 в гладкомышечных клетках пузирного протока**).
5. Увеличение сопротивления общего желчного протока (**повышенная экспрессия ЦОГ-2 в гладкомышечных клетках сфинктера Одди**).

Механизм развития патологических нарушений

Повышенная экспрессия ЦОГ-2 в гладкомышечных клетках стенки желчного пузыря способствует снижению эвакуаторной функции желчного пузыря и **“активного” пассажа печеночной желчи** в желчный пузырь (рис. 21).




Рис. 21. “Активный” и “пассивный” пассаж печеночной желчи в желчный пузырь и двенадцатiperстную кишку у больных хроническим некалькулезным холециститом с билиарным сладжем.

1 – печеночная желчь;
2 – пузирная желчь с билиарным сладжем.

Избыточная экспрессия ЦОГ-2 в эпителиальных клетках стенки желчного пузыря способствует снижению абсорбционной функции желчного пузыря (уменьшение абсорбции воды и билиарного холестерина в фосфолипидных везикулах) и **“пассивного” пассажа печеночной желчи** в желчный пузырь. Это сопровождается уменьшением концентрации общих желчных кислот в пузирной желчи и увеличением концентрации билиарного холестерина в фосфолипидных везикулах и способствует нарушению коллоидной стабильности пузирной




Рис. 21а. Пассаж печеночной желчи в желчный пузырь и в просвет ДПК у больных хроническим некалькулезным холециститом с билиарным сладжем, недостаточностью сфинктера Одди и дуоденогастральным желчным рефлюксом (хронический желчный гастрит). 1 = неконцентрированная печеночная желчь; 2 = слабо концентрированная пузирная желчь.




Рис. 21б. Пассаж печеночной желчи в желчный пузырь и в просвет ДПК у больных хроническим некалькулезным холециститом с билиарным сладжем и III билиарным типом дисфункции сфинктера Одди (гипертонус сфинктера общего желчного протока). 1 = неконцентрированная печеночная желчь; 2 = слабо концентрированная пузирная желчь.




Рис. 21в. Пассаж печеночной желчи в желчный пузырь и в просвет ДПК у больных хроническим некалькулезным холециститом с билиарным сладжем и III панкреатическим типом дисфункции сфинктера Одди (хронический спастический панкреатит). 1 = неконцентрированная печеночная желчь; 2 = слабо концентрированная пузирная желчь.




Рис. 21г. Пассивный пассаж печеночной желчи в желчный пузырь и сока поджелудочной железы в просвет ДПК у больных хроническим некалькулезным холециститом с билиарным сладжем и билиарно-панкреатическим рефлюксом (хронический билиарный панкреатит). 1 = неконцентрированная печеночная желчь; 2 = слабо концентрированная пузырная желчь.




Рис. 21д. Пассаж печеночной желчи в желчный пузырь и в просвет ДПК у больных хроническим некалькулезным холециститом с билиарным сладжем и синдромом избыточного бактериального роста в тонкой кишке (повышение давления в ДПК). 1 = неконцентрированная печеночная желчь; 2 = слабо концентрированная пузырная желчь.




Рис. 21е. Пассаж печеночной желчи в желчный пузырь и в просвет ДПК у больных хроническим некалькулезным холециститом с билиарным сладжем после лечения препаратами целекоксив и урсодезоксихолевой кислоты (нормальная функция сфинктера Одди). 1 = неконцентрированная печеночная желчь; 2 = слабо концентрированная пузырная желчь.

желчи и преципитации гранул билирубината кальция и кристаллов моногидрата холестерина, т.е. формированию “литогенной” пузырной желчи (рис. 22).



Рис. 22. Механизм формирования литогенной пузырной желчи у больных хроническим некалькулезным холециститом с билиарным сладжем.

Повышенная экспрессия ЦОГ-2 в эпителиальных клетках стенки желчного пузыря способствует гиперсекреции гликопротеинового муцина в просвет желчного пузыря и пузырную желчь. Увеличение концентрации гликопротеинового муцина свыше 2 мг/мл способствует его полимеризации и повышению вязкости пузырной желчи. Преципитация гранул билирубината кальция и кристаллов моногидрата холестерина в полимеризованных нитях гликопротеинового муцина способствует формированию билиарного сладжа, повышению его эхогенности и его “проявлению” при ультрасонографическом исследовании.

Снижение “активного” и “пассивного” пассажа печеночной желчи в желчный пузырь способствует увеличению пассажа печеночной желчи в двенадцатерстную кишку и пузырно-независимой энтерогепатической циркуляции желчных кислот, билиарного холестерина и билирубина (рис. 23).




Рис. 23. Энтерогепатическая циркуляция желчных кислот у больных хроническим некалькулезным холециститом с билиарным сладжем. 1 – пузырно-зависимая энтерогепатическая циркуляция желчных кислот; 2 – пузырно-независимая энтерогепатическая циркуляция желчных кислот; 3 – поступление желчных кислот в печень по печеночной артерии; 4 – синтез холевой кислоты; 5 – синтез хенодезоксихолевой кислоты; 6 – поступление желчных кислот в печень по воротной вене. **ПА** – печеночная артерия; **ПВ** – печеночная вена; **ВВ** – воротная вена; **ЖК** – желчные кислоты.

Увеличению пузырно-независимой энтерогепатической циркуляции желчных кислот способствует увеличению концентрации желчных кислот в гепатоцитах и снижению накопи-

тельно-выделительной функции печени (т.е. формированию хронического "мягкого" внутрипеченочного холестаза) (рис. 23).




Рис. 24. Обмен холестерина у больных хроническим некалькулезным холециститом с билиарным сладжем. 1 – пузырно-зависимый выход билиарного холестерина; 2 – пузырно-независимый выход билиарного холестерина; 3 – пузырно-печеночная циркуляция абсорбированного билиарного холестерина; 4 – гидролиз эфиров холестерина; 5 – биосинтез холестерина; 6 – синтез эфиров холестерина; 7 – гидролиз эфиров холестерина. **ХсБ** – холестерин безводный; **ХсМ** – моногидрат холестерина; **ПА** – печеночная артерия; **ПВ** – печеночная вена; **ВВ** – воротная вена; **ЛС** – лимфатические сосуды.

Увеличению пузырно-независимой энтерогепатической циркуляции билиарного холестерина способствует увеличению абсорбции билиарного холестерина в тонкой кишке, поступлению билиарного холестерина в гепатоциты и повышенной секреции в печеночную желчь (рис. 24).

Эти два фактора способствуют формированию "литогенной" печеночной желчи (рис. 25).




Рис. 25. Механизм формирования литогенной печеночной желчи у больных хроническим некалькулезным холециститом с билиарным сладжем.

Снижение пузырно- зависимого выведения билиарного холестерина и уменьшение концентрации желчных кислот в пузырной желчи способствуют формированию "литогенной" пузырной желчи и преципитации кристаллов моногидрата холестерина в просвете желчного пузыря у 100% больных хроническим некалькулезным холециститом с билиарным сладжем (рис. 26).




Рис. 26. Обмен холестерина и желчных кислот у больных хроническим некалькулезным холециститом с билиарным сладжем. 1 – биосинтез холестерина; 2 – синтез эфиров холестерина; 3 – гидролиз эфиров холестерина; 4 – биосинтез желчных кислот. РХМ – ремнантные хиломикроны; ЖК – желчные кислоты; ХсБ – холестерин безводный; ХсМ – моно-гидрат холестерина; ПА – печеночная артерия; ПВ – печеночная вена; ВВ – воротная вена; ЛС – лимфатические суды.

Длительная персистенция билиарного сладжа и гипомоторная дисфункция желчного пузыря способствуют формированию желчных камней в желчном пузыре и трансформации хронического некалькулезного холецистита с билиарным сладжем в хронический калькулезный холецистит.

Патогенетическое лечение больных хроническим некалькулезным холециститом с билиарным сладжем

Соответственно, лечение хронического некалькулезного холецистита с билиарным сладжем (с болевым синдромом) и с целью профилактики хронического калькулезного холецистита, дуоденогастрального рефлюкса, атрофического гастрита антрального отдела желудка и хронического билиарного панкреатита:

1. Целекоксиб – по 100 мг 2 раза в день после еды – 5-7 дней, после чего
2. Урсодезоксихолевая кислота – по 750 мг 1 раз в день на ночь – 2 месяца.

Целекоксиб – селективный ингибитор ЦОГ-2, ингибируя активность ЦОГ-2 в гладкомышечных клетках стенки желчного пузыря и пузырного протока, способствует устранению болевого синдрома в течение 3-5 дней, восстановлению эвакуаторной функции желчного пузыря и пузырно-зависимого выведения билиарного холестерина, “активного” и “пассивного” пассажа печеночной желчи в желчный пузырь и снижению пузырно-независимой энтерогепатической циркуляции желчных кислот, билиарного холестерина и билирубина.

Целекоксиб – селективный ингибитор ЦОГ-2, ингибируя активность ЦОГ-2 в эпителиальных клетках стенки желчного пузыря, способствует снижению секреции гликопротеинового муцина в просвет желчного пузыря, концентрации гликопротеинового муцина в пузырной желчи и вязкости пузырной желчи, что предупреждает формирование билиарного сладжа.

Сниженная активность ЦОГ-2 в эпителиальных клетках стенки желчного пузыря помогает восстанавливать абсорбционную функцию желчного пузыря (абсорбцию воды и билиарного холестерина из фосфолипидных везикул, что способствует увеличению концентрации желчных кислот и уменьшению концентрации билиарного холестерина в пузырной желчи).

Урсодезоксихолевая кислота – гидрофильная гепатозащитная желчная кислота способствует растворению кристаллов моноциклического холестерина в желчном пузыре, снижению литогенности пузырной и печеночной желчи, разрешению хронического “мягкого” внутрипеченочного холестаза (т.е. способствует восстановлению накопительно-выделительной функции печени) (1-66).

Целекоксиб и **Урсодезоксихолевая кислота**, патогенетически блокируя основные механизмы образования желчных камней, способствуют элиминации билиарного сладжа в 100% случаях и снижают риск повторного образования билиарного сладжа и, соответственно, уменьшают риск образования желчных камней в желчном пузыре (1-66).

Целекоксиб – селективный ингибитор ЦОГ-2, ингибируя активность ЦОГ-2 в гладкомышечных клетках желчевыводящих протоков и сфинктера Одди, способствует устранению болевого синдрома в течение 3-5 дней и восстановлению пассажа печеночной желчи в две-

надцатиперстную кишку.

Целекоксиб – селективный ингибитор ЦОГ-2, ингибируя активность ЦОГ-2 в эпителиальных клетках желчевыводящих протоков, способствует снижению секреции гликопротеинового муцина в просвет желчевыводящих протоков, концентрации гликопротеинового муцина в печеночной желчи и вязкости печеночной желчи, что предупреждает формирование билиарного сладжа и желчных камней в желчевыводящих протоках.

Сниженная активность ЦОГ-2 в эпителиальных и гладкомышечных клетках желчевыводящих протоков помогает уменьшить риск развития холедохолитиаза.

Урсодезоксихолевая кислота – гидрофильная гепатозащитная желчная кислота способствует растворению кристаллов моногидрата холестерина в желчевыводящих протоках, снижению литогенности печеночной желчи, разрешению хронического “мягкого” внутрипеченочного холестаза (т.е. способствует восстановлению накопительно-выделительной функции печени) и у части больных способствует растворению билиарного сладжа в желчевыводящих протоках (1-66).

Урсодезоксихолевая кислота – гидрофильная гепатозащитная желчная кислота, снижая агрессивные свойства желчи, препятствует развитию хронического атрофического гастрита антрального отдела желудка (дуоденогастральный рефлюкс, реактивный гастрит или желчный гастрит), хронического билиарного панкреатита (билиопанкреатический рефлюкс) и хронического спасищеского панкреатита (III панкреатический тип дисфункции сфинктера Одди) (1-66).

Целекоксиб и Урсодезоксихолевая кислота, патогенетически блокируя основные механизмы образования желчных камней, способствуют профилактике образования желчных камней в желчевыводящих протоках и снижают риск развития холедохолитиаза и билиарного панкреатита (1-66).

Эффективность – 95%.

Продолжительность ремиссии – 18-24 месяца.

Внимание!!! Информация для больных:

Перед использованием данной схемы лечения убедительная просьба посмотреть противопоказания (см. ниже) и побочные эффекты при применении фармакологических препаратов Целекоксиба и Урсодезоксихолевой кислоты и получить разрешение у вашего лечащего врача.

Противопоказания для применения Целекоксиба:

- аллергические реакции (крапивница, бронхоспазм) на прием ацетилсалициловой кислоты или других НПВС (в анамнезе);
- III триместр беременности;
- известная повышенная чувствительность к сульфонамидам;
- повышенная чувствительность к любому компоненту препарата.

Противопоказания для применения Урсодезоксихолевой кислоты:

- повышенная чувствительность к препарату;
- острые воспалительные заболевания желчного пузыря и желчных протоков;
- неспецифический язвенный колит;
- болезнь Крона.

Сайт не несет юридической ответственности за использование представленных схем лечения без согласования с лечащим врачом.

Список литературы

1. Mackay S, Dillane P. Biliary pain. *Aust Fam Physician*. 2004; **33(12)**: 977-981.
2. Lee SP, Nicholls JF. Nature and composition of biliary sludge. *Gastroenterology*. 1986; **90(3)**: 677-686.
3. Carey MC, Cahalane MJ. Whither biliary sludge? *Gastroenterology*. 1988; **95(2)**: 508-523.
4. Wilkinson LS, Levine TS, Smith D, Chadwick SJ. Biliary sludge: can ultrasound reliably detect the presence of crystals in bile? *Europ J Gastroenterol Hepatol*. 1996; **8(10)**: 999-1001.
5. Inoue K, Fuchigami A, Higashide S, Sumi S, Kogure M, Suzuki T, Tobe T. Gallbladder sludge and stone formation in relation to contractile function after gastrectomy. *Ann Surg*. 1992; **215(1)**: 19-26.

6. **Jüngst D**, Del Pozo R, Christoph S, Miquel JF, Eder MI, Lange V, Frimberger E, Von Ritter C, Paumgartner G. Quantification of biliary sludge in patients with cholesterol, mixed and pigment stones. *Gastroenterology*. 1994; **106(4)**: Abstr. 912.
7. **Jüngst D**, Del Pozo R, Christoph S, Miquel JF, Eder MI, Lange V, Frimberger E, Von Ritter C, Paumgartner G. Sedimentation of biliary sludge. Effect on composition of gallbladder bile from patients with cholesterol, mixed and pigment stones. *Scand J Gastroenterol*. 1996; **31(3)**: 273-278.
8. **Jüngst D**, Del Pozo R, Dolu MH, Schneeweiss SG, Frimberger E. Rapid formation of cholesterol crystals in gallbladder bile is associated with stone recurrence after laparoscopic cholecystotomy. *Hepatology*. 1997; **25(3)**: 509-513.
9. **Jüngst D**, Niemeyer A, Müller I, Zündt B, Meyer G, Wilhelm M, Del Pozo R. Mucin and phospholipids determine viscosity of gallbladder bile in patients with gallstones. *World J Gastroenterol*. 2001; **7(2)**: 203-207.
10. **Gründel D**, Jüngst C, Straub G, Althaus R, Schneider B, Spelsberg FW, Hüttl TP, Del Pozo R, Jüngst D, Neubrand M. Relation of gallbladder motility to viscosity and composition of gallbladder bile in patients with cholesterol gallstones. *Digestion*. 2009; **79(4)**: 229-234.
11. **Ilchenko AA**, Delyukina O.V. Clinical role of biliary sludge. *Consilium Medicum: Gastroenterology*. 2005; 2.
12. **Bilhartz LE**, Horton JD. Gallstone disease and its complications. In: M. Feldman, B.F. Scharschmidt, M.H. Sleisenger, eds. *Sleisenger and Fordtran's Gastrointestinal and Liver Disease: Pathophysiology, Diagnosis, Management*. 6th ed. Philadelphia: WB Saunders Company, 1998: 948-972.
13. **Pazzi P**, Petroni ML, Prandini N, Adam JA, Gullini S, Northfield TC, Jazrawi RP. Postprandial refilling and turnover: specific gallbladder motor function defects in patients with gallstone recurrence. *Eur J Gastroenterol Hepatol*. 2000; **12(7)**: 787-794.
14. **Jazrawi RP**, Pazzi P, Petroni ML, Prandini N, Paul C, Adam JA, Gullini S, Northfield TC. Postprandial gallbladder motor function: refilling and turnover of bile in health and in cholelithiasis. *Gastroenterology*. 1995; **109(2)**: 582-591.
15. **Svanvik J**, Thornell E, Zettergren L. Gallbladder function in experimental cholecystitis. *Surgery*. 1981; **89(4)**: 500-506.
16. **Minetoma T**. Relationship between gallbladder contractility and muscular fibrosis in the patients with cholecystolithiasis – immunohistochemical analysis. *Nippon Shokakibyo Gakkai Zasshi*. 1993; **90(12)**: 3018-3027.
17. **Hidaka T**, Nakano M, Inokuchi T, Sugiyama M, Nishi J, Ogura R. Arachidonate metabolism in bovine gallbladder mucosa. *Kurume Med J*. 1991; **38(3)**: 129-133.
18. **Nilsson B**, Delbro D, Hedin L, Friman S, Andius S, Svanvik J. Role of cyclooxygenase-2 for fluid secretion by the inflamed gallbladder mucosa. *J Gastrointest Surg*. 1998; **2(3)**: 269-277.
19. **Chapman WC**, Peterkin GA, LaMorte WW, Williams LF Jr. Alterations in biliary motility correlate with increased gallbladder prostaglandin synthesis in early cholelithiasis in prairie dog. *Dig Dis Sci*. 1989; **34(9)**: 1420-1424.
20. **MacPherson BR**, Pemsingh RS. Ground squirrel model for cholelithiasis: role of epithelial glycoproteins. *Microsc Res Tech*. 1997; **39(1)**: 39-55.
21. **Pemsingh RS**, Macpherson BR, Scott GW. Characterization of lipid accumulation in the gallbladder mucosa of the Ground Squirrels fed a lithogenic diet. *J Pathology*. 1988; **154(2)**: 173-180.
22. **Pemsingh RS**, Macpherson BR, Scott GW. Mucus hypersecretion in the gallbladder epithelium of Ground Squirrels fed a lithogenic diet for the induction of cholesterol gallstones. *Hepatology*. 1987; **7(6)**: 1267-1271.
23. **Pemsingh RS**, Macpherson BR, Scott GW. Morphological observations on the gallbladder of Ground Squirrels fed a lithogenic diet. *J Pathology*. 1987; **152(2)**: 127-135.
24. **Macpherson BR**, Pemsingh RS, Scott GW. Experimental cholelithiasis in the Ground Squirrel. *Laboratory Investigation*. 1987; **56(2)**: 138-145.
25. **Sasaki H**, Tazuma S, Kajiyama G. Effects of 16,16-dimethyl prostaglandin E2 on biliary mucous glycoprotein and gallstone formation in guinea pigs. *Scand J Gastroenterol*. 1993; **28(6)**: 495-499.
26. **Meyers D**, Feldstein DA. Initial treatment of biliary colic: are NSAIDs better than opiates? *WJM*. 2005; **104(4)**: 9.
27. **Myers SI**, Bartula LL, Colvin MP, Parkman HP. Cholecystokinin (CCK) down regulates PGE2 and PG12 release in inflamed Guinea pig gallbladder smooth muscle cell cultures. *Prostaglandins Leukot Essent Fatty Acids*. 2005; **73(2)**: 121-126.
28. **Myers SI**, Riva A, Kalley-Taylor B, Bartula L. Taurodeoxycholic acid stimulates rabbit gallbladder eicosanoid release. *Prostaglandins Leukot Essent Fatty Acids*. 1995; **52(1)**: 35-39.
29. **Myers SI**, Inman LR, Kalley-Taylor B, Riva A, Bartula L. Increased intragallbladder pressure stimulates gallbladder eicosanoid release. *Prostaglandins*. 1994; **48(1)**: 53-66.
30. **Myers SI**, Bartula L. Human cholecystitis is associated with increased gallbladder prostaglandin I2 and prostaglandin E2 synthesis. *Hepatology*. 1992; **16(5)**: 1176-1179.
31. **Myers SI**, Bartula LL. Sex differences in gallbladder prostaglandin synthesis mediated by acute inflammation. *Prostaglandins Leukot Essent Fatty Acids*. 1990; **41(4)**: 259-264.
32. **Kaminski DL**. Arachidonic acid metabolites in hepatobiliary physiology and disease. *Gastroenterology*. 1989; **97(3)**: 781-792.
33. **Kaminski DL**, Deshpande Y, Thomas L, Qualy J, Blank W. Effect of oral ibuprofen on formation of prostaglandins E and F by human gallbladder muscle and mucosa. *Dig Dis Sci*. 1985, **30(10)**: 933-940.
34. **Xiao ZL**, Amaral J, Biancani P, Behar J. Impaired cytoprotective function of muscle in human gallbladders with cholesterol stones. *Am J Physiol Gastrointest Liver Physiol*. 2005; **288(3)**: G525-G532.

35. Prystowsky JB, Rege RV. The inflammatory effects of crystalline cholesterol monohydrate in the guinea pig gallbladder *in vivo*. *Surgery*. 1998; **123**(3): 258-263.
36. Torsoli A, Corazziari E, Habib FI, Cicala M. Pressure relationships within the human bile tract. Normal and abnormal physiology. *Scand J Gastroenterol Suppl*. 1990; **175**: 52-57.
37. Shoda J, Ueda T, Kawamoto T, Todoroki T, Asano T, Sugimoto Y, Ichikawa A, Maruyama T, Nimura Y, Tanaka N. Prostaglandin E receptors in bile ducts of hepatolithiasis patients and the pathobiological significance for cholangitis. *Clin Gastroenterol Hepatol*. 2003; **1**(4): 285-96.
38. Kano M, Shoda J, Satoh S, Kobayashi M, Matsuzaki Y, Abei M, Tanaka N. Increased expression of gallbladder cholecystokinin: a receptor in prairie dogs fed a high-cholesterol diet and its dissociation with decreased contractility in response to cholecystokinin. *J Lab Clin Med*. 2002; **139**(5): 85-94.
39. Shoda J, Kano M, Asano T, Irimura T, Ueda T, Iwasaki R, Furukawa M, Kamiya J, Nimura Y, Todoroki T, Matsuzaki Y, Tanaka N. Secretory low-molecular-weight phospholipases A2 and their specific receptor in bile ducts of patients with intrahepatic calculi: factors of chronic proliferative cholangitis. *Hepatology*. 1999; **29**(4): 1026-1036.
40. Kano M, Shoda J, Irimura T, Ueda T, Iwasaki R, Urasaki T, Kawauchi Y, Asano T, Matsuzaki Y, Tanaka N. Effects of long-term ursodeoxycholate administration on expression levels of secretory low-molecular-weight phospholipases A2 and mucin genes in gallbladders and biliary composition in patients with multiple cholesterol stones. *Hepatology*. 1998; **28**(2): 302-313.
41. Shoda J, Ueda T, Ikegami T, Matsuzaki Y, Satoh S, Kano M, Matsuura K, Tanaka N. Increased biliary group II phospholipase A2 and altered gallbladder bile in patients with multiple cholesterol stones. *Gastroenterology*. 1997; **112**(6): 2036-2047.
42. Krishnamurthy S, Krishnamurthy GT. Biliary dyskinesia: role of the sphincter of Oddi, gallbladder and cholecystokinin. *J Nucl Med*. 1997; **38**(11): 1824-1830.
43. Isogai M, Yamaguchi A, Hori A, Nakano S. Hepatic histopathological changes in biliary pancreatitis. *Amer J Gastroenterol*. 1995; **90**(3): 449-454.
44. Honda A, Yoshida T, Tanaka N, Matsuzaki Y, He B, Shoda J, Osuga T. Increased bile acid concentration in liver tissue with cholesterol gallstone disease. *J Gastroenterol*. 1995; **30**(1): 61-66.
45. Geraghty JM, Goldin RD. Liver changes associated with cholecystitis. *J Clin Pathol*. 1994; **47**(5): 457-460.

Список литературы

- Целекоксиб (Celecoxib) и Урсодезоксихолевая кислота (ursodeoxycholic acid):
1. Chen XW, Cai JT. The impact of selective cyclooxygenase-2 inhibitor celecoxib on the formation of cholesterol gallstone. *Zhonghua Nei Ke Za Zhi*. 2003; **42**(11): 797-799.
 2. Joshi GP. Valdecoxib for the management of chronic and acute pain. *Expert Rev Neurother*. 2005; **5**(1): 11-24.
 3. Jayr C. Analgesic effects of cyclooxygenase 2 inhibitors. *Bull Cancer*. 2004; **91** (Suppl 2): S125-S131.
 4. Kumar A, Deed JS, Bhasin B, Kumar A, Thomas S. Comparison of the effect of diclofenac with hyoscine-N-butylbromide in the symptomatic treatment of acute biliary colic. *ANZ J Surg*. 2004; **74**(7): 573-576.
 5. Matheson AJ, Figgitt DP. Rofecoxib: a review of its use in the management of osteoarthritis, acute pain and rheumatoid arthritis. *Drugs*. 2001; **61**(6): 833-865.
 6. Akriavidis EA, Hatzigavriel M, Kapnias D, Kirimlidis J, Markantas A, Garyfallos A. Treatment of biliary colic with diclofenac: a randomized, double-blind, placebo-controlled study. *Gastroenterology*. 1997; **113**(1): 225-231.
 7. Añez MS, Martínez D, Pacheco JL, González H, Rivera J, Pelaschier E, Uzcátegui L, Romero MD, Molina Z, Roditti de Montilla M, et al. Indomethacin in the treatment of acute cholecystitis and biliary colic. *G E N*. 1991; **45**(1): 32-37.
 8. Goldman G, Kahn PJ, Alon R, Wiznitzer T. Biliary colic treatment and acute cholecystitis prevention by prostaglandin inhibitor. *Dig Dis Sci*. 1989; **34**(6): 809-811.
 9. Kaminski DL, Deshpande Y, Thomas L, Qualy J, Blank W. Effect of oral ibuprofen on formation of prostaglandins E and F by human gallbladder muscle and mucosa. *Dig Dis Sci*. 1985; **30**(10): 933-940.
 10. Ikegami T, Matsuzaki Y, Fukushima S, Shoda J, Olivier JL, Bouscarel B, Tanaka N. Suppressive effect of ursodeoxycholic acid on type IIA phospholipase A2 expression in HepG2 cells. *Hepatology*. 2005; **41**(4): 896-905.
 11. Kano M, Shoda J, Irimura T, Ueda T, Iwasaki R, Urasaki T, Kawauchi Y, Asano T, Matsuzaki Y, Tanaka N. Effects of long-term ursodeoxycholate administration on expression levels of secretory low-molecular-weight phospholipases A2 and mucin genes in gallbladders and biliary composition in patients with multiple cholesterol stones. *Hepatology*. 1998; **28**(2): 302-313.
 12. Guarino MP, Carotti S, Morini S, Perrone G, Behar J, Altomare A, Alloni R, Caviglia R, Emerenziani S, Rabitti C, Cicala M. Decreased number of activated macrophages in gallbladder muscle layer of cholesterol gallstone patients following ursodeoxycholic acid. *Gut*. 2008; **57**(12): 1740-1741.
 13. Carotti S, Guarino MP, Cicala M, Perrone G, Alloni R, Segreto F, Rabitti C, Morini S. Effect of ursodeoxycholic acid on inflammatory infiltrate in gallbladder muscle of cholesterol gallstone patients. *Neurogastroenterol Motil*. 2010; **22**(8): 866-873.
 14. Mizuno S, Tazuma S, Kajiyama G. Stabilization of biliary lipid particles by ursodeoxycholic acid. Prolonged nucleation time in human gallbladder bile. *Dig Dis Sci*. 1993; **38**(4): 684-693.
 15. Tazuma S, Sasaki H, Mizuno S, Sagawa H, Hashiba S, Horiuchi I, Kajiyama G. Effect of ursodeoxy-

- cholic acid administration on nucleation time in human gallbladder bile. *Gastroenterology*. 1989; **97**(1): 173-178.
16. **Jüngst C**, Sreejayan N, Zündt B, Müller I, Spelsberg FW, Hüttl TP, Kullak-Ublick GA, del Pozo R, Jüngst D, von Ritter C. Ursodeoxycholic acid reduces lipid peroxidation and mucin secretagogue activity in gallbladder bile of patients with cholesterol gallstones. *Eur J Clin Invest*. 2008; **38**(9): 634-639.
17. **Fischer S**, Müller I, Zündt BZ, Jüngst C, Meyer G, Jüngst D. Ursodeoxycholic acid decreases viscosity and sedimentable fractions of gallbladder bile in patients with cholesterol gallstones. *Eur J Gastroenterol Hepatol*. 2004; **16**(3): 305-311.
18. **Sauter GH**, Thiessen K, Parhofer KG, Jüngst C, Fischer S, Jüngst D. Effects of ursodeoxycholic acid on synthesis of cholesterol and bile acids in healthy subjects. *Digestion*. 2004; **70**(2): 79-83.
19. **Fahey DA**, Carey MC, Donovan JM. Bile acid/phosphatidylcholine interactions in mixed monomolecular layers: differences in condensation effects but not interfacial orientation between hydrophobic and hydrophilic bile acid species. *Biochemistry*. 1995; **34**(34): 10886-10897.
20. **Guarino MP**, Carotti S, Sarzano M, Alloni R, Vanni M, Grossi M, Sironi G, Maffettone PL, Cicala M. Short-term ursodeoxycholic acid treatment improves gallbladder bile turnover in gallstone patients: a randomized trial. *Neurogastroenterol Motil*. 2005; **17**(5): 680-686.
21. **Guarino MP**, Cong P, Cicala M, Alloni R, Carotti S, Behar J. Ursodeoxycholic acid improves muscle contractility and inflammation in symptomatic gallbladders with cholesterol gallstones. *Gut*. 2007; **56**(6): 815-820.
22. **Mas MR**, Comert B, Mas N, Yamaneli L, Ozotuk H, Tasci I, Jazrawi RP. Effects of long term hydrophilic bile acid therapy on in vitro contraction of gallbladder muscle strips in patients with cholesterol gallstones. *World J Gastroenterol*. 2007; **13**(32): 4336-4339.
23. **Colecchia A**, Mazzella G, Sandri L, Azzaroli F, Magliuolo M, Simoni P, Bacchi-Reggiani ML, Roda E, Festi D. Ursodeoxycholic acid improves gastrointestinal motility defects in gallstone patients. *World J Gastroenterol*. 2006; **12**(33): 5336-5343.
24. **Xiao ZL**, Biancani P, Carey MC, Behar J. Hydrophilic but not hydrophobic bile acids prevent gallbladder muscle dysfunction in acute cholecystitis. *Hepatology*. 2003; **37**(6): 1442-1450.
25. **van de Heijning BJ**, van de Meenberg PC, Portincasa P, Doornewaard H, Hoebers FJ, van Erpecum KJ, Vanberge-Henegouwen GP. Effects of ursodeoxycholic acid therapy on in vitro gallbladder contractility in patients with cholesterol gallstones. *Dig Dis Sci*. 1999; **44**(1): 190-196.
26. **Mendez-Sanchez N**, Brink MA, Paigen B, Carey MC. Ursodeoxycholic acid and cholesterol induce enterohepatic cycling of bilirubin in rodents. *Gastroenterology*. 1998; **115**(3): 722-732.
27. **Beuers U**. Drug insight: Mechanisms and sites of action of ursodeoxycholic acid in cholestasis. *Nat Clin Pract Gastroenterol Hepatol*. 2006; **3**(6): 318-328.
28. **Pemberton PW**, Aboutwerat A, Smith A, Warnes TW. Ursodeoxycholic acid in primary biliary cirrhosis improves glutathione status but fails to reduce lipid peroxidation. *Redox Rep*. 2006; **11**(3): 117-123.
29. **Jeong HJ**, Kim CG. Pretreatment with ursodeoxycholic acid (UDCA) as a novel pharmacological intervention in hepatobiliary scintigraphy. *Yonsei Med J*. 2005; **46**(3): 394-398.
30. **Lukivskaya OY**, Maskevich AA, Buko VU. Effect of ursodeoxycholic acid on prostaglandin metabolism and microsomal membranes in alcoholic fatty liver. *Alcohol*. 2001; **25**(2): 99-105.
31. **Bouscarel B**, Ceryak S, Robins SJ, Fromm H. Studies on the mechanism of the ursodeoxycholic acid-induced increase in hepatic low-density lipoprotein binding. *Lipids*. 1995; **30**(7): 607-617.
32. **Bomzon A**, Ljubuncic P. Ursodeoxycholic acid and in vitro vasoactivity of hydrophobic bile acids. *Dig Dis Sci*. 2001; **46**(9): 2017-2024.
33. **Ljubuncic P**, Said O, Ehrlich Y, Meddings JB, Shaffer EA, Bomzon A. On the in vitro vasoactivity of bile acids. *Br J Pharmacol*. 2000; **131**(3): 387-398.
34. **Sinisalo J**, Vanhanen H, Pajunen P, Vapaatalo H, Nieminen MS. Ursodeoxycholic acid and endothelial-dependent, nitric oxide-independent vasodilatation of forearm resistance arteries in patients with coronary heart disease. *Br J Clin Pharmacol*. 1999; **47**(6): 661-665.
35. **Pak JM**, Adeagbo AS, Triggle CR, Shaffer EA, Lee SS. Mechanism of bile salt vasoactivity: dependence on calcium channels in vascular smooth muscle. *Br J Pharmacol*. 1994; **112**(4): 1209-1215.
36. **Otake M**, Sandoh N, Sakaguchi T, Tsukada K, Hatakeyama K. Enhancement of portal blood flow by ursodeoxycholic acid in partially hepatectomized rats. *Surg Today*. 1996; **26**(2): 142-144.
37. **Bomzon A**, Ljubuncic P. Bile acids as endogenous vasodilators? *Biochem Pharmacol*. 1995; **49**(5): 581-589.
38. **Pak JM**, Lee SS. Vasoactive effects of bile salts in cirrhotic rats: in vivo and in vitro studies. *Hepatology*. 1993; **18**(5): 1175-1181.
39. **Benedetti A**, Alvaro D, Bassotti C, Gigliozi A, Ferretti G, La Rosa T, Di Sario A, Baiocchi L, Jezequel AM. Cytotoxicity of bile salts against biliary epithelium: a study in isolated bile ductule fragments and isolated perfused rat liver. *Hepatology*. 1997; **26**(1): 9-21.
40. **Itoh S**, Kono M, Akimoto T. Psoriasis treated with ursodeoxycholic acid: three case reports. *Clin Exp Dermatol*. 2007; **32**(4): 398-400.
41. **Günsar C**, Melek M, Karaca I, Sencan A, Mir E, Ortaç R, Canan O. The biochemical and histopathological effects of ursodeoxycholic acid and metronidazole on total parenteral nutrition-associated hepatic dysfunction: an experimental study. *Hepatogastroenterology*. 2002; **49**(44): 497-500.
42. **Tomida S**, Abei M, Yamaguchi T, Matsuzaki Y, Shoda J, Tanaka N, Osuga T. Long-term ursodeoxycholic acid therapy is associated with reduced risk of biliary pain and acute cholecystitis in patients with gallbladder stones: a cohort analysis. *Hepatology*. 1999; **30**(1): 6-13.
43. **Okoro N**, Patel A, Goldstein M, Narahari N, Cai Q. Ursodeoxycholic acid treatment for patients with postcholecystectomy pain and bile microlithiasis. *Gastrointest Endosc*. 2008; **68**(1): 69-74.
44. **Guma C**, Viola L, Apéstegui C, Pinchuk L, Groppa J, Michelini J, Martínez B, Bolaños R, Toselli L.

- Therapeutic efficacy of ursodeoxycholic acid in persistent gallbladder lithiasis and persistent biliary sludge: preliminary results of a multicenter experience. *Acta Gastroenterol Latinoam.* 1994; **24(4)**: 233-237.
45. **Ros E**, Navarro S, Bru C, Garcia-Pugés A, Valderrama R. Occult microlithiasis in «idiopathic» acute pancreatitis: prevention of relapses by cholecystectomy or ursodeoxycholic acid therapy. *Gastroenterology.* 1991; **101(6)**: 1701-1709.
46. **Testoni PA**, Caporuscio S, Bagnolo F, Lella F. Idiopathic recurrent pancreatitis: long-term results after ERCP, endoscopic sphincterotomy, or ursodeoxycholic acid treatment. *Am J Gastroenterol.* 2000; **95(7)**: 1702-1707.
47. **Borda F**, Oquiñena S, Borobio E, Vila J, Frauca A, Martínez B. Is pre-operative treatment with ursodeoxycholic acid useful in reducing relapses in acute biliary pancreatitis? *An Sist Sanit Navar.* 2003; **26(2)**: 225-229.
48. **Okazaki K**. Therapy for chronic pancreatitis and the prognosis. *Nihon Naika Gakkai Zasshi.* 2004; **93(1)**: 45-50.
49. **Saraswat VA**, Sharma BC, Agarwal DK, Kumar R, Negi TS, Tandon RK. Biliary microlithiasis in patients with idiopathic acute pancreatitis and unexplained biliary pain: response to therapy. *J Gastroenterol Hepatol.* 2004; **19(10)**: 1206-1211.
50. **Venneman NG**, van Berge-Henegouwen GP, van Erpecum KJ. Pharmacological manipulation of biliary water and lipids: potential consequences for prevention of acute biliary pancreatitis. *Curr Drug Targets Immune Endocr Metabol Disord.* 2005; **5(2)**: 193-198.
51. **Venneman NG**, van Erpecum KJ. Gallstone disease: Primary and secondary prevention. *Best Pract Res Clin Gastroenterol.* 2006; **20(6)**: 1063-1073.
52. **Tsubakio K**, Kiriyama K, Matsushima N, Taniguchi M, Shizusawa T, Katoh T, Manabe N, Yabu M, Kanayama Y, Himeno S. Autoimmune pancreatitis successfully treated with ursodeoxycholic acid. *Intern Med.* 2002; **41(12)**: 1142-1146.
53. **Okazaki K**. Ursodeoxycholic acid as an alternative therapy for autoimmune pancreatitis. *Intern Med.* 2002; **41(12)**: 1082-1083.
54. **Scarpa PJ**, Cappell MS. Treatment with ursodeoxycholic acid of bile reflux gastritis after cholecystectomy. *J Clin Gastroenterol* 1991; **13(5)**: 601-603.
55. **Realini S**, Reiner M, Frigerio G. Treatment of dyspeptic disorders, lithiasis and biliary dyskinesia with ursodeoxycholic acid. Analysis of a controlled multicenter study. *Schweiz Med Wochenschr.* 1980; **110(22)**: 879-880.
56. **Alvisi V**, Tralli M, Loponte A, D'Ambrosi A, Pavani F, Ruina M. Ursodeoxycholic acid in the treatment of dyspeptic-painful disorders of biliary origin: report of a controlled multicenter study. *Clin Ter.* 1982; **100(1)**: 21-33.
57. **Stefaniwsky AB**, Tint GS, Speck J, Shefer S, Salen G. Ursodeoxycholic acid treatment of bile reflux gastritis. *Gastroenterology* 1985; **89(5)**: 1000-1004.
58. **Pazzi P**, Stabellini G. Effect of ursodeoxycholic acid (UDCA) on biliary dyspepsia in patients without gallstones. *Cur Ther Res* 1985; **37**: 685-690.
59. **Rosman AS**. Efficacy of ursodeoxycholic acid (UDCA) in treating bile reflux gastritis. *Gastroenterology.* 1987; **92(1)**: 269-272.
60. **Scalia S**, Pazzi P, Stabellini G, Guarneri M. HPLC assay of conjugated bile acids in gastric juice during ursodeoxycholic acid (Deursil) therapy of bile reflux gastritis. *J Pharm Biomed Anal.* 1988; **6(6-8)**: 911-917.
61. **Pazzi P**, Scalia S, Stabellini G. Bile reflux gastritis in patients without prior gastric surgery: Therapeutic effects of ursodeoxycholic acid. *Cur Ther Res* 1989; **45**: 476-680.
62. **Scarpa PJ**, Cappell MS, Chen WY, Liao WC. Treatment with ursodeoxycholic acid of bile reflux gastritis after cholecystectomy. *J Clin Gastroenterol.* 1991; **13(5)**: 601-603.
63. **Mathai E**, Arora A, Cafferkey M, Keane CT, O'Morain C. The effect of bile acids on the growth and adherence of *Helicobacter pylori*. *Aliment Pharmacol Ther.* 1991; **5(6)**: 653-658.
64. **Piepoli AL**, Caroppo R, Armentano R, Caruso ML, Guerra V, Maselli MA. Tauroursodeoxycholic acid reduces damaging effects of taurodeoxycholic acid on fundus gastric mucosa. *Arch Physiol Biochem.* 2002; **110(3)**: 197-202.
65. **Ozkaya M**, Erten A, Sahin I, Engin B, Ciftci A, Cakal E, Caydere M, Demirbaş B, Ustün H. The effect of ursodeoxycholic acid treatment on epidermal growth factor in patients with bile reflux gastritis. *Turk J Gastroenterol* 2002; **13(4)**: 198-202.
66. **Thao TD**, Ryu HC, Yoo SH, Rhee DK. Antibacterial and anti-atrophic effects of a highly soluble, acid stable ursodeoxycholic acid (UDCA) formula in *Helicobacter pylori*-induced gastritis. *Biochem Pharmacol.* 2008; **75(11)**: 2135-2146.